



## A NUTRITIOUS AND ECO-FRIENDLY CROP: MUSHROOMS

Salman Khan<sup>1</sup>, Tanvi Tripathi<sup>1</sup>, Shikha Tripathi<sup>1</sup>, Vivek Kumar<sup>1</sup>, Sharukh Hasan<sup>1</sup>,  
Saurabh Dubey<sup>1</sup> & Avinash Pratap Singh<sup>2</sup>

<sup>1</sup>Research Scholar & <sup>2</sup>Assistant Professor

Dept. of Botany, CMP Degree College, University of Allahabad, Prayagraj

E-mail: avinashsinghau@gmail.com

### Abstract

Mushrooms are an extremely nutritious and environmentally friendly crop with several medical benefits. Mushrooms have less protein than mammals, while plants have far more. Except for iron, it has a greater concentration of fibre, vital amino acids, and minerals. It is not only a nutrient-rich vegetable high in vitamin D, but it can also prevent cancer, HIV-1, AIDS (a terrible human disease), and other illnesses. It is a crop that requires minimal resources and area to develop and is cultivated worldwide at a low cost throughout the year. Growing mushrooms from agricultural waste substrates is a cost-effective and environmentally friendly method for producing nutritious and delicious food. Lignocellulosic waste can also be converted into food, feed, and fertilizers through cultivation. Mushroom consumption and production are low compared to other crops, resulting in limited investment in the business.

**Keywords:** Edible Mushrooms, Food Industry, Medicinal Properties, Nutritive Value.

### Introduction

Mushrooms have long been farmed for their nutritional benefits and flavour, particularly in eastern nations. Edible mushroom farming is crucial in today's world, especially with a growing population and environmental challenges. This research highlighted key mushroom species farmed in India, their nutritional and medicinal benefits, and environmentally sustainable production practices. Cultivated mushrooms are a very nutritious food that may be cultivated from biological, agricultural, or agro-industrial waste (Sánchez, 2010; Atila, 2017). Researchers found nutritional differences between mushrooms grown on various substrates. Hoa et al. (2015) found that formulas containing 100% sugar bagasse and 100% corncob had higher levels of protein, fiber, ash, and mineral content (Ca, K, and Mg) compared to 100% sawdust. Supplementing *Agaricus bisporus* mushroom substrates with trace elements has been shown to produce fruiting bodies rich in micronutrients like selenium, copper, and zinc, which are often lacking in human diets. Bird et al. (2017); Rzymiski et al. (2017); Werner and Beelman (2002). Mushrooms are used to make a variety of nutritional goods. Mushrooms are valued for their antiviral, anti-cancerous, and antioxidant effects, leading to increased interest in their production and processing. Edible mushrooms are widely consumed due to their high protein, vitamins, minerals, fiber, microelements, and low-calorie content (Naeem et al., 2020). Universal cultivated mushrooms. Commercial cultivation of mushrooms is limited to a few kinds of fungi. Many of the 300 genera of mushrooms and associated fleshy basidiomycetes do not sporulate without a host due to mycorrhizal conditions. However, many saprophytic plants are cultivable.

### Nutritional Value of Mushrooms

Edible mushrooms provide a nutrient-dense and readily digested diet. Mushrooms are used for their nutritional benefits, flavour, and taste. According to Aremu et al. (2009), mushrooms have more protein

content than fruits and vegetables. Mushrooms have a similar dry weight to yeast and are heavier than dried beans and peas. Nutritional requirements vary among species and substrates. According to Koyyalamudi et al. (2009), mushrooms are beneficial for heart patients due to their high protein and low-calorie content, as well as their ability to provide all essential amino acids. Tryptophan and lysine have higher concentrations than cysteine and methionine. Edible mushrooms are low in carbohydrates and fat. Mushrooms are an ideal meal for diabetics and people looking to lose weight, owing to their low starch content.

According to Haas and James (2009), edible mushrooms can also be considered a vegetable meat. This versatile ingredient may enhance the flavour of other recipes. Some vegetarians augment their diet with mushrooms instead of meat (Abulude, 2005).

Researchers Kattawan et al. (2011) and Naeem et al. (2020) found that mushrooms contain antioxidants. Mushrooms have potential medical advantages, including anticancer properties, in addition to their nutritional value. Several scholars (Abulude, 2005; Kuforiji and Fasidi, 2008; Kattawan et al., 2011) have discussed the therapeutic properties of *Pleurotus tuber-regium*.

Mushrooms may be used with medicinal herbs to treat a variety of ailments, including chest discomfort, headache, dropsy, fever, smallpox, and stomach pain (Change, 2012). Mushrooms have several nutritional benefits, similar to those found in cereals, legumes, and meat. According to Patel and Goyal (2013), mushrooms are low in calories, fat, salt, and cholesterol.

Mushrooms have anti-cancer properties, acting as an immediate anti-mitotic, oxygen species inducer, mitotic kinase inhibitor, topoisomerase inhibitor, and angiogenesis inhibitor. This leads to apoptosis and cancer dispersal.

### **Medicinal value**

Mushrooms contain a variety of chemical substances that may have medicinal properties. *Grifola*'s lipid fraction has antioxidant properties and inhibits enzymes linked to chronic illnesses, including cancer (Inoue et al., 2002). According to Chang (1996), mushroom ingredients not only use direct cytotoxicity against tumour cells to prevent disease progression, but also block non-immune processes.

*Ganoderma* species produce compounds with antibacterial properties (Smania et al., 1999; Mothana et al., 2000), inhibiting the development of bacteria such as *Staphylococcus*. Kuznetsov et al. (2005) found that steroid isolates had antimicrobial properties. Mushrooms have been shown to have nutritional and therapeutic benefits for ages, with increasing experimental data supporting these claims.

Medicinal mushroom experiments for HIV/AIDS patients in Africa have shown good results (Chang, 2006), increasing their worth. According to Badalyan (2004), mycelial extracts of *Lentinula* have antiprotozoal properties against *Paramecium*. Mushrooms exhibit antiviral capabilities, with isolated components from *Ganoderma* being effective against HIV-1 (El-Mekkawy et al., 1998; Ichimura et al., 1998). They also have antiviral effects against the influenza virus type 1.

Mushroom extracts are utilized for medical purposes and as therapies. *Lentinula edodes* (shiitake), *Grifola frondosa* (maitake), and *Ganoderma lucidum* (reishi) have been used medicinally for millennia in Asia. Medicinal mushrooms have been studied for their antiparasitic, cardiovascular, anticancer, antibacterial, antiviral, anti-inflammatory, antidiabetic, and hepatoprotective effects (Lentinan, 2009). According to Akpaja (2005) and Smith et al. (2002), mushrooms have medical, nutritional, and mythical functions. Mushroom genetic resources are widely used in various industries, including agriculture, agronomy, animal feed, and human food. They can also be used to produce, develop, and detect high-value constituents in industries such as pharmaceuticals and chemicals. Mushroom compounds, such as polysaccharides, proteoglycans, and glycoproteins, can regulate immune responses and inhibit tumour development. Extracts are often used in China, Japan, and Korea to supplement chemotherapy and radiation treatments (Smith et al., 2002; Borchers et al., 2008).

Mushrooms containing psychoactive chemicals have been utilized for therapeutic purposes, both mental and physical (Huder 2000). Sarfaraz et al. (2009) suggest that some mushrooms, particularly polypores like Reishi, can assist a number of health conditions. A study by Ogbe et al. (2008) found that using *Ganoderma* sp. improved avian egg laying capability and illness resistance. Okhuoya et al. (2010) discovered that a diet based on Betaglucan from *Ganoderma lucidum* (Leyss.) can effectively treat buruli ulcers caused by *Mycobacterium* in Nigeria.

### Other Uses of Mushrooms

Mushrooms are used to colour wood and other natural fibers. Mushroom dyes organic chromophores generate brilliant and solid hues, allowing for a wide range of colours. According to Mussak and Bechtold (2009), natural dyes were formerly used as a source for many colours. In industrialized nations, mushrooms have been utilized as a fire starter. Evocative Device LLC uses them for biodegradable packaging. They contribute significantly to the advancement of innovative biological filtration and remediation methods. Wikipedia (accessed in 2011). According to Akpaja et al. (2005), mushrooms may also be used as cannon powder.

### Environment-friendly crop

The use of an expensive substrate for growing oyster mushrooms increases their cost of production. There is a need to search for definite substitute materials that should exist in sufficient amounts at a relatively low price (Arya and Arya, 2003). *Pleurotus* has been reported to grow readily on many non-conventional substrates (Das et al., 2000; Mukherjee and Nandi, 2002; Nageswaran et al., 2003). In West Bengal, India, water hyacinth (*Eichhornia crassipes* Solms.), a low-cost supplement for oyster mushroom (*Pleurotus florida*) cultivation. Bandopadhyay et al. (2009). In India, *Pleurotus sajor-caju* has been successfully cultivated on banana pseudo-stem and paddy straw (Jandaik, 1974; Jandaik and Kapoor, 1975).

Jandaik and Kapoor (1975) conducted experiments on rice straw, wheat straw, ragi straw, hulled maize cob, and waste paper. Unused garbage and rubbish may be used to grow protein-rich mushrooms and reduce environmental pollution. (Medda 2001). Biotechnological processes may recycle large amounts of organic waste through mushroom culture, producing food, fuel, and fertilizers. Using spent mushroom cultivation leftovers as a biologically pre-treated substrate might improve biogas production. Agro waste is recycled through mushroom growing (Madan, 1994).

### Conclusion

As the world's population grows, the supply of food and medical services decreases, particularly for people in developing nations. Mushrooms are a cost-effective and environmentally friendly source of nourishment and medicine, with several species available.

Mushrooms may grow on many waste materials from agriculture and industry, such as paper, sawdust, cardboard, and wood. Mushroom substrates, after harvesting, are utilized in agricultural cultivation as compost. Mushroom growing is a cost-effective, nutritious, therapeutic, and environmentally friendly crop that also recycles waste materials.

### Conflict of Interest

There is no conflict of interest

### References

1. Abulude, F. O. (2005). Proximate and phytate compositions of mushrooms consumed in South Western Nigeria. *Advances in Food Science*, 27(4), 185–188.
2. Akpaja, E. O., Okhuoya, J. A., & Heferere, B. A. E. (2005). Ethnomycology and indigenous cases of mushrooms among the Bini speaking people of Nigeria: A case study of aihuobabekun community near Benin City, Nigeria. *International Journal of Medicinal Mushroom*, 7(3), 373–374.
3. Aremu, M. O., Basuk, Gyan, S. D., Goyal, A., Bhowmik, P. K., & Banik, S. D. (2009). Proximate composition and functional properties of mushroom flours from *Ganoderma* spp., *Omphalotus olearius* (DC) Sing, and

- Hebeloma mesphaeum* (Pers) Quels used in Nassarawa State, Nigeria. *Mal of Journal Nutrition*, 15(2), 233–241.
4. Arya, C., & Arya, A. (2003). Effect of acid hydrolysis of substrate on yield of oyster mushroom *Pleurotus sajor-caju* (Fr.) Singer. *Mushroom Res*, 12, 35–38.
  5. Atila, F. (2017). Evaluation of suitability of various agro-wastes for productivity of *Pleurotus djamor*, *Pleurotus citrinopileatus* and *Pleurotus eryngii* mushrooms. *J. Exp. Agric. Int.*, 17(5), 1–11.
  6. Badalyan, S. M. (2004). Antiprotozoal activity and mitogenic effect of mycelium of culinary medicinal shiitake mushroom *Lentinus edodes* (Berk.) Singer (Agaricomycetidae). *Int. J. Med. Mushrooms*, 6, 131–138.
  7. Bandopadhyay, S., Khatun, S., Mitra, S., Roy, P., Dasgupta, A., Chaudhuri, S. K., & Chattopadhyay, N. C. (2009). Antihyperglycaemic effect of dietary mushroom (*Pleurotus florida*) in alloxan-induced diabetic rats. In *Proc. 5th Int. Medicinal Mushroom Conference*, Mycological Society of China, Nantong, China (pp. 135–141).
  8. Bird, J. K., Murphy, R. A., Ciappio, E. D., & McBurney, M. I. (2017). Risk of deficiency in multiple concurrent micronutrients in children and adults in the United States. *Nutrients*, 9(7), 655.
  9. Borchers, A. T., Krishnamurthy, A., Keen, C. L., Meyers, F. J., & Gershwin, M. E. (2008). The immunobiology of mushrooms. *Experimental Biology and Medicine*, 233(93), 259–276.
  10. Chang, S. T. (1996). Mushroom research and development—equality and mutual benefit. In *Proceedings of the 2nd International Conference on Mushroom Biology and Mushroom Products*, Pennsylvania State University, Pennsylvania, USA (pp. 1–10).
  11. Chang, S. T. (2006). The world mushroom industry: Trends and technological developments. *Int. J. Med. Mushrooms*, 8, 297–314.
  12. Change, R. (2012). Functional properties of edible mushrooms. *Nutrition Reviews*, 54, 91–93.
  13. Das, N., Mahapatra, S. C., & Chattopadhyay, R. N. (2000). Use of wild grasses as substrate for cultivation of oyster mushrooms in South West Bengal. *Mushroom Res*, 9, 95–99.
  14. El-Mekkawy, S., Meselhy, M. R., Nakamura, N., Tezuka, Y., Hattori, M., & Kakiuchi, N. (1998). Anti-HIV-1 and anti-HIV-1 protease substances from *Ganoderma lucidum*. *Phytochem*, 49, 1651–1657.
  15. Haas, E. M., & James, P. (2009). *More vegetables, please!! Delicious recipes for eating healthy food each and every day*. Oakland, CA: New Harbinger Publications. pp. 222. ISBN 97815-72245907.
  16. Hoa, H. T., Wang, C. L., & Wang, C. H. (2015). The effects of different substrates on the growth, yield and nutritional composition of two oyster mushrooms (*Pleurotus ostreatus* and *Pleurotus cystidiosus*). *Mycobiology*, 43(4), 423–434.
  17. Huder, G. W. (2000). *Magical Mushrooms, Mischievous Mold*. Princeton, NJ: Princeton University Press. pp. 175. ISBN 0-691-07016-4.
  18. Ichimura, T., Watanabe, O., & Muruyama, S. (1998). Inhibition of HIV-1 protease by water-soluble lignin-like substance from an edible mushroom, *Fuscoporia obliqua*. *Biosci. Biotechnol. Biochem*, 62, 575–577.
  19. Inoue, A., Kodama, N., & Nanba, H. (2002). Effect of maitake (*Grifola frondosa*) D-fraction on the control of the T lymph node Th-1/Th-2 proportion. *Biol. Pharm. Bull*, 25, 536–540.
  20. Jandaik, C. L., & Kapoor, J. N. (1975). Cultural studies on some edible fungi. *Indian J. Mushrooms*, 1, 22–26.
  21. Jandaik, C. L. (1974). Artificial cultivation of the mushroom *Pleurotus sajor-caju* (Fr.) Singer. *Mushroom J*, 22, 405.
  22. Kattawan, A., Chanlekha, K., Kongkachuichai, R., & Chaaroensiri, R. (2011). Effects of cooking on antioxidant activities and polyphenol content of edible mushrooms commonly consumed in Thailand. *Pakistan Journal of Nutrition*, 10(11), 1094–1103.
  23. Koyyalamudi, S. R., Jeong, S. C., Song, C. H., Cho, K. Y., & Pang, G. (2009). Vitamin D2 for the nation and bioavailability from *Agaricus bisporus* button mushrooms treated with ultraviolet irradiation. *Journal of Agriculture and Food Chemistry*, 57(8), 3351–3355.
  24. Kuforiji, O., & Fasidi, I. (2008). Compositional studies on *Pleurotus tuber-regium* sclerotia. *Advances in Food Sciences*, 30(1), 2–5.

25. Kuznetsov, O. I., Milkova, E. V., Sosnia, A. E., & Sotnikova, N. I. (2005). Antimicrobial action of *Lentinus edodes* juice on human microflora. *Mikrobiol. Epidem. Immunobiol*, 1, 80–82.
26. Labarère, J., & Menini, G. U. (2000). Collection, characterization, conservation and utilization of mushrooms, germplasm resources in Africa. In *Proceedings of the First International Congress for the Characterization, Conservation, Evaluation and Utilization of Mushroom Genetic Resources for Food and Agriculture*. FAO, Bordeaux, France, 9–13.
27. Lentinan. (2009). About herbs. Memorial Sloan Kettering Cancer Center. Madan, M. (1994). Mushroom cultivation for rural development. In *Souvenir National Symposium on Mushrooms* (Solan, H. P., Eds.), pp. 82–87.
28. Medda, R. N. (2001). *Studies on nutritional requirements for the cultivation of edible mushrooms and the associated yield* (Ph.D. thesis, Department of Botany, Burdwan University, West Bengal, India), 6 pp.
29. Mothana, R. A. A., Jansen, R., Julich, W. D., & Lindequist, U. (2000). Ganomycin A and B, new antimicrobial farnesyl hydroquinones from the basidiomycete *Ganoderma pfefferi*. *J. Nat. Prod*, 63, 416–418.
30. Mukherjee, R., & Nandi, B. (2002). Role of nutrient supplementation on productivity of *Pleurotus* spp. on two lignocellulosic biomass and dry matter digestibility of the spent substrate. In N. Samajpati (Ed.), *Tropical Mycology: Proc. of Third Nat. Symposium* (Indian Mycol. Soc., Kolkata, pp. 180–188).
31. Mussak, R., & Bechtold, T. (2009). *Handbook of natural colorants*. New York: Wiley. pp. 183–200. ISBN 0-470-51199-0.
32. Naeem, M. Y., Ugur, S., & Rani, S. (2020). Emerging role of edible mushrooms in food industry and its nutritional and medicinal consequences. *Eurasian Journal of Food Science and Technology*, 4(1), 6–23.
33. Nageswaran, M., Gopalakrishnan, M., Ganesan, M., Vedhamurthy, A., & Selaganapadhyay, E. (2003). Evaluation of water hyacinth for culture of oyster mushroom. *J. Aqua. Plant Manag*, 41, 122–123.
34. Ogbe, A. O., Mgbogikwe, L. O., Owoade, A. A., Atawodi, S. E., & Abdu, A. P. (2008). The effect of wild mushrooms (*Ganoderma lucidum*) supplementation of seed on the immune response of pullet chickens to infections, bursal disease cancer. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 7(4), 2844–2855.
35. Okhuoya, J. A., Akpaja, E. O., Osemwegie, O. O., Ogherekano, A. O., & Ihayere, C. A. (2010). Nigerian mushrooms: Underutilized non-wood forest resource. *Journal of Applied Science and Environmental Management*, 14(1), 43–54.
36. Osemwegie, O. O., Eriyaremu, E. G., & Abdulmahik, J. (2006). A survey of macrofungi in Edo/Delta region of Nigeria, their morphology and uses. *Global Journal of Pure and Applied Science*, 12(2), 149–157.
37. Patel, S., & Goyal, A. (2013). Recent developments in mushrooms as anti-cancer therapeutics. *Journal of Biotechnology*, 2, 1–15.
38. Rzymiski, P., Mleczek, M., Niedzielski, P., Siwulski, M., & Gąsecka, M. (2017). Cultivation of *Agaricus bisporus* enriched with selenium, zinc and copper. *J. Sci Food Agric*, 97(3), 923–928.
39. Sánchez, C. (2010). Cultivation of *Pleurotus ostreatus* and other edible mushrooms. *Appl. Microbiol. Biotechnol*, 85, 1321–1337.
40. Sarfaraz, K. W., Mir, A. K., Muhammad, A. K., Mushtaq, A., Muhammed, Z., Fazal-Ur-Rehman, & Shazia, S. (2009). Vegetables mentioned in the Holy Quran and Ahadith and their ethnomedical studies in Dera Ismail Khan N.W.F.P., Pakistan. *Pakistan Journal of Nutrition*, 8(5), 530–538.
41. Smania, A., Delle, M. F., Smania, E. F. A., & Cuneo, R. S. (1999). Antibacterial activity of steroidal compounds isolated from *Ganoderma applanatum* (Pers.) Pat. (Aphyllphoromycetidae) fruit body. *International Journal of Medicinal Mushrooms*, 1, 325–330.
42. Smith, J. E., Rowan, N. J., & Sullivan, R. (2002). Medicinal mushrooms: A rapidly developing area of biotechnology for cancer therapy and other bioactivities. *Biotechnol. Lett*, 24, 1839.
43. Werner, A. R., & Beelman, R. B. (2002). Growing high-selenium edible and medicinal button mushrooms [*Agaricus bisporus* (J. Lge) Imbach] as ingredients for functional foods or dietary supplements. *Int. J. Med. Mushrooms*, 4, 88–94.